In the Specification

Please replace the paragraph beginning at line 8, on page 1 with the following rewritten paragraph:

This application is a continuation of U.S. Application Serial No. 09/272,097, filed on March 18, 1999, which is a continuation of U.S. Application Serial No. 09/046,203, filed on March 23, 1998, now Patent No. 5,945,526, which is a continuation of U.S. Application No. 08/726,462, filed on October 4, 1996, now Patent No. 5,800,996, which is a continuation-in-part of U.S. Application No. 08/672,196, filed on June 27, 2996, now U.S. Patent No. 5,847,162 which is a continuation-in-part of U.S. Application No. 08/642,330, filed on May 3, 1996, now Patent No. 5,863,727, each of which it is incorporated herein by reference in its entirety.

Please replace the paragraph beginning at line 20, on page 10, with the following rewritten paragraph:

In another embodiment, the energy transfer fluorescent dyes have donor and acceptor dyes with the general structure where Y_1 and Y_2 taken separately are either hydroxyl, oxygen, iminium or amine, the iminium and amine preferably being a tertiary iminium or amine and R_{11} - R_{16} are any substituents which are compatible with the energy transfer dyes of the present invention. /

please replace all of page 52 with the following rewritten paragraphs:

In compound 3A-A, one of R_1 and R_2 is ethyl, the other being hydrogen, R_3 and R_4 taken separately are hydrogen, R_6 is methyl, R_5 and R_7 - R_{10} taken separately are hydrogen, X_1 is carboxylate, and one of X_3 and X_4 is a linking group, the other being hydrogen.

In compound 3A-B, one of R_1 and R_2 is ethyl, the other being hydrogen, R_3 and R_4 taken separately are methyl, R_5 is methyl, R_6 - R_{10} taken separately are hydrogen, X_1 is carboxylate, and, one of X_3 and X_4 is a linking group, the other being hydrogen.

In compound 3A-C, R_1 and R_2 taken separately are methyl, R_3 and R_9 taken together form a six membered ring, R_4 and R_8 taken together form a six membered ring, R_5 , R_6 , R_7 , and R_{10} taken separately are hydrogen, X_1 is carboxylate, and, one of X_3 and X_4 is a linking group, the other being hydrogen.

In compound 3B-D, R_1 and R_2 taken separately are hydrogen, R_3 and R_9 taken together form a six membered ring, R_4 and R_8 taken together form a six membered ring, R_5 , R_6 , $R_{7,a}$ and R_{10} taken separately are hydrogen, X_1 is carboxylate, and one of X_3 and X_4 is a linking group, the other being hydrogen.

In compound 3B-E, one of R_1 and R_2 is ethyl, the other being hydrogen, R_3 and R_9 taken together form a six membered ring, R_4 and R_8 taken together form a six membered ring, R_5 is methyl, R_6 , R_7 and R_{10} taken separately are hydrogen, X_1 is carboxylate, and, one of X_3 and X_4 is a linking group, the other being hydrogen.

In compound 3B-F, R_1 and R_2 taken separately are hydrogen, R_3 and R_4 taken separately are methyl, R_5 - R_{10} taken separately are hydrogen, X_1 is carboxylate, and, one of X_3 and X_4 is linking group, the other being hydrogen.

Please replace the paragraph beginning at line 5, on page 53 with the following rewritten paragraph:

Figure 4A shows a generalized synthesis wherein the substituent X_1 can be other than carboxylate. In the figure, X' indicates moieties which are precursors to X_1 . In the method illustrated in Figure 4A, two equivalents of a 3-aminophenol derivative 4A-A/4A-B, such as 3-dimethylaminophenol, is reacted with one equivalent of a dichlorobenzene derivative 4A-C, e.g., 4-carboxy-3,6,dichloro-2-sulfobenzoic acid cyclic anhydride, i.e., where the X_1' moieties of 4c taken together are,

JPlease replace the paragraphs beginning at line14, on page 53 with the following rewritten paragraphs:



The reactants are then heated for 12 h in a strong acid, e.g., polyphosphoric acid or sulfuric acid, at 180°C. The crude dye 4A-D is precipitated by addition to water and isolated